




## A study of relapse rate and functional outcome after surgical debridement in patients with chronic osteomyelitis

Masthan Basha Shaik , Kiran Kumar Mallam , Divya Vurundhur , Anil Babu Talari 

Department of Orthopaedics, ACSR Govt Medical College, Nellore, Andhra Pradesh, India

\* Correspondence: Masthan Basha Shaik. Department of Orthopaedics, ACSR Govt Medical College, Nellore, Andhra Pradesh, India. Tel: +919494585868, Email: [bonesbasha@gmail.com](mailto:bonesbasha@gmail.com)

### Abstract

**Background:** This study analyzed the rate of relapse and functional outcome after surgery for chronic osteomyelitis of long bones.

**Methods:** This study included 100 cases of chronic osteomyelitis, which were treated with surgical debridement, along with a 6-week course of antibiotics. The patients were divided into two groups based on whether they had undergone preoperative magnetic resonance imaging (MRI) or not. A comparison of relapse rates was then conducted between these two groups. The patients were followed upto 9<sup>th</sup> month postop, and their functional outcomes were examined using the Association for the Study and Application of the Method of Ilizarov Criteria (ASAMI), Lower Extremity Functional Scale (LEFS), Disabilities of the Arm, Shoulder, and Hand (DASH), and Musculoskeletal Tumor Society (MSTS) questionnaires. Logistic regression analysis was used to assess the impact of risk factors on the recurrence or relapse.

**Results:** Sixty patients had chronic osteomyelitis, and 40 patients were infected non-union. Twenty-five patients had preoperative MRI. The mean preoperative LEFS was  $44 \pm 13.59$  (19-73). Preoperative ASAMI showed 4% excellent, 62% fair, 24% good, and 10% were poor. The preoperative MSTS emotional component showed that enthusiastic (0%), satisfied (0%), accepts (36%), and dislikes (60%). Among the infected non-union cases, 2 cases had a total disability, 15 cases had recreational restriction, and 23 cases had a partial disability. Majority cases were gram-positive organism cultured were methicillin-sensitive *Staphylococcus aureus* accounts 21%, and methicillin-resistant *S. aureus* accounts 16%. Common antibiotics administered were Cloxacillin in 23 cases, Meropenem in 11 cases, Septran in 8 cases, Linezolid in 8 cases, and Vancomycin in 7 cases. The mean duration of antibiotics administered was 6 weeks. At 9<sup>th</sup> month postoperative, ASAMI showed a reduction in the number of patients categorized as 'fair' from 12 to 9, while the number of patients classified as 'good' increased from 2 to 6 cases. Fourteen (66.6%) patients who were initially classified under the 'fair' or 'poor' category preoperatively decreased to 9 (42.8%) postoperatively. The number of patients in the 'good' category increased from 6 (28%) preoperatively to 9 (42.9%) at the 6-month postoperative mark. Of the 61 patients, 42 (64.6%) had significant functional improvement at 9<sup>th</sup> month postoperative among those who had chronic osteomyelitis, while of the 34 patients, 15 (44.1%) patients with infected non-union had significant functional improvement ( $P = 0.04$ ). 20.7% out of 77 patients who were culture positive went on to have relapse within 9<sup>th</sup> month follow-up and 15% out of the 20 patients who had no-growth on intraoperative cultures went on to develop relapse following 9<sup>th</sup> month post-surgery follow-up. Nine patients (24.3%) in the posttraumatic group had recurrence of infection, while 1 patient had relapse at 6 months following surgery. At the 6-month follow-up, the percentage of relapse was 13.3% ( $n = 2$ ) in the MRI group and 25% ( $n = 8$ ) in the non-MRI group.

**Conclusion:** The relapse rate was 19.8%. In total, 19.3% of surgical debridement had experienced a recurrence of infection within the 9 months follow-up, and 7.3% of patients required repeat surgery during the hospital stay. Factors that determined recurrence were age > 55 years, long bone involvement, fracture, monotherapy, receiving < 6 weeks of antibiotics, and inadequate treatment. Disease burden, financial conflicts, disability, and long duration of treatment affect the patient's emotional status. These factors may also affect the patient's compliance with the prescribed treatment plan.

### Article History

Received: 01 November 2023  
Received in revised form: 4 December 2023  
Accepted: 10 December 2023  
Published online: 15 January 2024  
DOI: [10.29252/JCBR.7.4.10](https://doi.org/10.29252/JCBR.7.4.10)

### Keywords

Osteomyelitis  
Debridement  
Emotional status  
Functional outcome  
Relapse rate  
Musculoskeletal Tumour Society

Article Type: Original Article



### Highlights

#### What is current knowledge?

- The emotional and physical strain on individuals with chronic osteomyelitis had a tremendous impact.
- The complex course of the osteomyelitis disease, high recurrence rates, and the treatment involved pose a heavy burden on osteomyelitis patients.

#### What is new here?

- The study assessed various causative organisms, duration of antibiotics administered, and functional outcomes in patients with chronic osteomyelitis who are undergoing surgical debridement.
- Measured the relapse rates in different age groups of patients with chronic osteomyelitis.
- The study compares the relapse rates in osteomyelitis patients with/without preoperative MRI.

### Introduction

Chronic osteomyelitis is caused by untreated or incompletely treated hematogenous osteomyelitis, untreated open fractures, or surgical infection complications. Recurrence with the same organism isolate is referred to as "relapse," while recurrence with a different or extra strain is referred to as "re-infection" (1).

There has also been a continual evolution in diagnostic techniques, novel antibiotics, biomaterials, and reconstructive operations (2).

Antibiotic therapy alone will not suffice due to its poor kinetics in bone with sequestrae and fistulae. Currently, surgical debridement is the preferred method, with minimum or no systemic antibiotics and/or local antibiotics (3,4).

The most difficult aspect of treating chronic osteomyelitis is that infection eradication is a major concern. Various patients relapse over time, necessitating several operations and lengthy hospitalizations, imposing a significant financial burden on patients and health care systems (5). Chronic osteomyelitis had a recurrence rate ranging from 0% to 48% (6,7,8). The emotional, physical, and financial burdens associated with chronic osteomyelitis have had a profound impact on the lives of millions of individuals worldwide (9).

A significant number of studies and new research continue to be conducted in the diagnosis and management of osteomyelitis. There is very little research on the functional result and relapse rates after surgery.

The primary objective of this study was to fill a knowledge gap by investigating the rate of relapse after surgical intervention. Additionally, the study aimed to evaluate the functional outcome of patients who underwent surgical debridement. Furthermore, the researchers aimed to describe the rate of recurrence or relapse of chronic osteomyelitis following surgical debridement and compare the functional outcomes during regular follow-up visits.

## Methods

Patients who were diagnosed with chronic osteomyelitis based on clinical, radiological, or culture evidence and underwent surgical debridement were included in the study.

Chronic osteomyelitis-host type A and B (Cierny and Mader) patients were studied.

The study was carried out in the Department of Orthopaedics at ACSR Govt Medical College in Nellore, Andhra Pradesh.

All patients with single long bone osteomyelitis or infected non-union, adults over the age of 18yrs, patients who had surgical debridement, and patients with a minimum follow-up of 3 months were included (Figure 1).

Exclusion criteria were patients with multifocal osteomyelitis, joint involvement, vascular compromise, Type C host, and bone loss (lengthening/bone transport).




Figure 1. Flowchart for selection criteria and study protocol

## Procedure

Patients were divided into groups based on whether or not a preoperative magnetic resonance imaging (MRI) was performed, and relapse rates were compared. Data were gathered from patients suffering from chronic osteomyelitis of the long bones. The Association for the Study and Application of the Method of Ilizarov Criteria (ASAMI), Musculoskeletal Tumor Society (MSTS-functional category), and Lower Extremity Functional Scale (LEFS) for lower limb long bone osteomyelitis, and Disabilities of the Arm, Shoulder, and Hand (DASH) questionnaires for upper limb long bone osteomyelitis were used to track patients' functional outcomes.

## Diagnosis

The relapse of chronic osteomyelitis was described as an infection arising again at the same spot from which it was previously thought to be cleared and which was specifically treated with surgery. On presentation, the infection is described as the growth of the same pathogen or a new pathogen in repeat cultures.

## Follow-up

A patient was termed relapse-free if there were no clinical symptoms of infection such as warmth, discharging sinus, fever, edema, or soreness during follow-up. The erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP) were used to confirm normalization. Participants were followed up of minimum of

3 months and maximum upto 9 months from February 2023 to October 2023. Study duration is Preoperative MRI or infected non-union/chronic osteomyelitis classified patients into groups.

The relapse is defined as the recurrence of infection in the same location where it was previously thought to be cleared and was particularly treated with surgery.

## Statistical analysis

Demographic, clinical, and laboratory characteristics were all documented in a proforma. An analysis of variance (ANOVA) was performed to analyze continuous variables with preoperative frequency data. Chi-square tests were used to examine categorical variables that had a functional consequence. Non-categorical results were compared using a Student's t-test. The relationship between the score and clinical profile was assessed using Pearson correlation analysis. P values less than 0.05 were considered statistically significant. Statistical analysis was done using SPSS version 22.0.

## Results

Among the 100 patients, 60 had chronic osteomyelitis, and 40 had infected non-union. The average age was 35 years. Males made up 82% of the patients.

The majority of patients (82%) were diagnosed with chronic osteomyelitis as a result of a posttraumatic origin, with hematogenous etiology accounting for 18%.

The long bones of the lower limb, such as the femur and tibia, were involved in 94 cases, whereas the humerus was involved in 6.

The majority of patients (45%) were semi-skilled, followed by 20% skilled, 15% shopkeepers, and so on.

Twelve individuals had diabetes, and 10 had hypertension.

Among walking aids, 35% of the patients were found to be using walking aids such as a crutch or walking stick before the index surgery.

To avoid inter-observer bias, an orthopedic surgeon validated the existence of sequestrum in 65 of 100 patients based on preoperative radiographs. Preoperative MRI was performed on 25 patients at the discretion of the surgeon.

## Functional outcome

### Lower Extremity Functional Scale functional score

The Lower Extremity Functional Scale consists of a series of 20 questions, each of which is rated on a 5-point ordinal scale ranging from 0 (very difficult/impossible to accomplish) to 4 (no difficulty). The functional capacity was to express nominally and was scored out of a maximum of 80, with a minimal observable change of 9 points regarded clinically significant. The mean preoperative LEFS was 44.1±3.59 (19 to 73).

### Musculoskeletal Tumor Society-emotional status

The emotional acceptance level was assessed at the time of presentation following a detailed explanation of the problem and the comprehensive management of the disease involved, including the possibility of recurrence, staged procedures, costs involved, several hospital visits, regular follow-up, and so on.

In our study, 60% of patients disliked their condition at the time of presentation, whereas 36% accepted it. In our study, the emotional aspect of MSTs indicates that none of the cases showed enthusiasm or satisfaction. However, 36% of cases were accepted, and 60% of cases were disliked.

### The association for the study and application of the method of Ilizarov criteria functional score

The ASAMI criteria contained 2 components: a radiologically measured bone score and a history and clinical examination-based functional score. The ASAMI functional score was divided into 4 categories as follows: Excellent: active, no limp, minimal stiffness (loss of 15° knee extension/dorsiflexion of ankles), no reflex sympathetic dystrophy, and no pain. Good: active with one or more of the following symptoms: limp, stiffness, Reflex Sympathetic Dystrophy (RSD), or considerable pain. Fair: active, with 3 or more of the following symptoms: limp, stiffness, RSD, and considerable pain. Poor: inactivity (unemployment or inability to do everyday activities due to injury) and amputation.

In our study, preoperative ASAMI was found to be outstanding (4%), fair (62%), good (24%), and bad (10%).

### Cierny-Mader classification (Physiological)

Most participants (83%) belong to host A, and 13% belong to host B.

### Cierny-Mader classification (Anatomical)

The majority were Cierny-made type IV patient population (42%), 9% were type I, 22% were type II, and 21% patients belonged to Cierny-mader host-A type.

### Musculoskeletal Tumor Society-Functional category

There are 22 cases of chronic osteomyelitis with partial impairment, 36 cases with recreational restriction, and 2 cases with no restriction. Among infected non-union cases, 2 had entire disability, 15 had recreational restrictions, and 23 had partial impairment.

## Culture reports

According to the culture reports, 78% of the samples were culture-positive. Methicillin-sensitive *Staphylococcus aureus* accounted for 21% of the gram-negative organisms grown, while methicillin-resistant *S. aureus* accounted for 16% of the samples (Table 1).

Table 1. The culture report of study samples

Gram-negative organism	Frequency %	Gram-positive organism	Frequency %
<i>Pseudomonas</i>	12	Methicillin-resistant <i>Staphylococcus aureus</i>	16
<i>Escherichia coli</i>	5	Methicillin-sensitive <i>Staphylococcus aureus</i>	21
Carbapenem-resistant organisms	6	Enterococcus	2
<i>Klebsiella</i>	7	B-hemolytic streptococci	3
Proteus	4		
NFGNB	2		
Enterobacter	1		

Abbreviation: NFGNB: Nonfermenting gram-negative bacilli

## Antibiotic treatment

Nine cases out of 19 with hematogenous osteomyelitis had intraoperative culture growth of methicillin-sensitive *S. aureus* and were given antibiotics for 6 weeks. In the intraoperative cultures of 4 patients, no growth was identified. Cloxacillin was given to 23 patients, Meropenem to 11 individuals, Septran to 8 patients, Linezolid to 8 patients, and Vancomycin to 7 patients. The average length of antibiotic administration was 6 5.06 weeks (95% CI, 4.3-6.4).

## Emotional status outcome

After 3 months, there was a decrease in the percentage of patients who disliked their condition (63% to 20%), as well as an increase in the percentage of patients who accepted and understood the problems with their illness and the weight of therapy (33% to 47%). Following surgery, 24% of patients were satisfied after 3 months, and none of the patients were excited at postoperative.

## Functional outcome

There was a total of 21 patients with infected non-union who returned for a 6-month follow-up. Preoperatively, 7 cases (33.4%) were in the partial disability category, while 5 cases (23.8%) were in the postoperative partial disability category. Recreational limitation was seen in 14 cases (67.6%), whereas the overall number of patients with recreational restriction after surgery was 16 (76.1%). There were no people who were completely disabled or had no limits. According to the MSTs-functional scale, the classification of the 17 patients with infected non-union 9<sup>th</sup> month postop is as follows:

## Preoperative vs 9<sup>th</sup> month postoperative MSTs-functional status in the infected non-union group

During the study period, 17 out of 29 patients (5 patients; ≤9months postoperative) were followed up 9<sup>th</sup> month after the index operation. The number of patients in the partial disability group decreased from 11 to 7, while the number of patients in the recreational limitation group increased from 5 to 10. In addition, 58.5% of patients with infected non-union were physically followed up for up to 9<sup>th</sup> month postop.

## 6-month postoperative change in ASAMI-functional outcome in patients with infected non-union and infected non-union group

The number of patients in the poor category decreased from 4 (16.6%) to 1 (4.1%). Preoperatively, there were 17 (70.7%) patients classified as fair, while postoperatively, the number decreased to 12 (50%) individuals. The number of patients classified as good increased from 3 (12.5%) preoperatively to 11 (45.8%) postoperatively. Notably, there was a significant improvement in functional outcomes among patients who progressed from the fair to good category and from the poor to fair category.

## Preoperative vs 9<sup>th</sup> month postoperative functional outcome infected non-union group

At 9<sup>th</sup> month postoperative, the number of patients in the fair group decreased from 12 to 9, whereas the number of patients in the good category increased from 2 to 6.

## Preoperative vs 6-month postoperative functional outcome in the chronic osteomyelitis group

There were 7 (33.2%) partially impaired patients preoperatively, while only 5 (22.8%) remained partially disabled postoperatively. At 6 months postoperatively, 2 patients in the partially impaired group who were classified as just recreationally restricted improved.

## Preoperative vs 6-month postoperative functional outcome (Based on ASAMI)

Preoperatively, 14 (66.7%) patients were in the fair or poor group; however, the

number of patients in this category decreased to 9 (42.8%) after surgery. The number of patients in the good group increased from 6 (28%) prior to surgery to 9 (42.9%) 6 months later, indicating an improvement in their functional status. According to Table 2, the number of patients in the excellent group increased from 1 (4.8%) preoperatively to 3 (14.3%) 6 months later.

Table 2. Preoperative vs 6-month postoperative functional outcome

Criteria	Functional outcome	Preoperative	6-month postoperative
ASAMI	Fair or less	14(66.7%)	9(42.9%)
	Good	6(28%)	9(42.9%)
	Excellent	1(4.8%)	3(14.3%)
MSTS	Total disability	0	0
	Partial disability	7(33.3%)	5(23.8%)
	Recreational restriction	14(66.7%)	16(76.2%)

Abbreviations: ASAMI, Association for the Study and Application of the Method of Ilizarov Criteria; MSTS, Musculoskeletal Tumor Society.

## Preoperative vs 9<sup>th</sup> month postop functional outcome (Based on MSTs-functional)

At 9<sup>th</sup> month postop follow-up, the number of partially disabled patients dropped from 6 (41.7%) preoperatively to 5 (35.72%). Preoperatively, there were 8 (58.3%) patients who had recreational restrictions, and this number decreased to 7 (50%) at the 9<sup>th</sup> month postop follow-up.

At 9<sup>th</sup> month follow-up, no restrictions were seen in 2 (14.28%) of cases (Table 3).

Table 3. Preoperative vs 9<sup>th</sup> month postop MSTs-functional outcome (n=14)

	Preoperative	9 <sup>th</sup> month postoperative
Total disability	0	0
Partial disability	6(41.7%)	5(35.72%)
Recreational restriction	8(58.3%)	7(50%)
No restriction	0	2(14.28%)

## Preoperative vs 9<sup>th</sup> month postoperative functional outcome (MSTS)

According to the functional outcome based on MSTS, 13 patients were classified as poor, fair, and good prior to surgery. Following surgery, the number of patients increased to 7 in the fair category, 5 in the good category, and 2 in the excellent category.

## Primary diagnosis vs clinical improvement at 9<sup>th</sup> month postop

At 9<sup>th</sup> month postop, 42 patients (64.6%) out of 61 patients with chronic osteomyelitis had substantial functional improvement, while 15 patients (44.1%) with infected non-union had significant functional improvement (P = 0.04; Table 4).

There was no substantial clinical improvement in 19 (42.2%) individuals with chronic osteomyelitis. This was related to a number of prognostic factors:

- Four patients underwent a primary flap procedure, after which they were immobilized for longer periods of time to minimize soft tissue issues that could harm the limb.
  - Seven patients had had more than 3 previous surgeries, either at our institute or elsewhere.
  - In intraoperative cultures, 5 individuals exhibited resistant *Pseudomonas aeruginosa*.
- Because 8 patients had concomitant joint stiffness prior to surgery, the prognosis was poor, and the breadth of functional recovery was limited.

Table 4. Association between diagnosis and clinical improvement at 9<sup>th</sup> month (LEFS)

Functional outcome (LEFS)	Chronic osteomyelitis (n=61)	Infected non-union (n=34)
Clinically improved	42(64.6%)	15(44.1%)
Not clinically improved	19(29.2%)	19(55.9%)

Abbreviation: LEFS, Lower Extremity Functional Scale.

The total number of patients examined at 6 months was 45, with 15.7% showing clinically meaningful improvement in the functional outcome, and the rest showed no improvement (Table 5).

One 9<sup>th</sup> month after surgery, there were 31 patients with clinical follow-ups, 17 of whom were infected non-union patients, and 14 were chronic osteomyelitis patients. According to LEFS, 15 patients (48.6%) showed significant functional improvement, whereas the remaining 16 (51.6%) showed no clinical improvement.

Table 5. Functional outcome at 6 months and 9<sup>th</sup> month postop (LEFS)

Clinically assessed functional outcome improvement		Frequency, n%
At 6 months	No clinically significant improvement	38 (84.3%)
	Clinically significant improvement	7(15.7%)
At 9 <sup>th</sup> month	No clinically significant improvement	16 (51.6%)
	Clinically significant improvement	15 (48.4%)



Relapse rates were examined among different patient groups based on criteria such as culture positivity, primary diagnosis, cause of osteomyelitis, and preoperative MRI.

Within 9<sup>th</sup> month postop follow-up, 20.78% of the 77 patients who were culture positive had recurrence, while 15% of the 20 patients who had no growth on intraoperative cultures had relapse.

There was a decrease in the number of patients who followed up at 6 months post-surgery for chronic osteomyelitis, with a total of 21 patients included in the follow-up. At 6 months post-surgery, 9 patients (24.3%) in the posttraumatic group had recurrence of infection, whereas 1 patient had relapse. At 6 months, the percentage of recurrence in the MRI group was 13.3% (n = 2) and 25% (n = 8) in the non-MRI group (Table 6).

**Table 6.** Association between culture positivity and relapse; primary diagnosis and relapse; preoperative MRI and relapse at 6 months

	Culture	No relapse	Relapse
Culture	+ve	61(79.12%)	16 (20.78%)
	-ve	17(85%)	3(15%)
Primary diagnosis	Chronic osteomyelitis	54(83.15%)	11(16.85%)
	Infected non-union	26(76.5%)	8(23.5%)
Preoperative MRI at 6 months	MRI	13(86.72%)	2(13.28%)
	Non-MRI	24(75%)	8(25%)

**Abbreviation:** MRI, magnetic resonance imaging.

## Discussion

Recurrence of chronic osteomyelitis has been observed even after 50 years of symptom-free interval. The recurring pattern of microbiological profiles is poorly known (10). The current study demonstrates a greater rate of infection recurrence or relapse in posttraumatic osteomyelitis. Chronic osteomyelitis relapsed at a rate of 19.8%.

According to the literature, the recurrence rate of chronic osteomyelitis ranged from 0% to 48%. According to Garcia del Pozo et al, 22.4% of people relapse (11).

Uçkay et al suggested that the bone once infected becomes a vulnerable area of bacterial adhesion when transient bacteremia occurs due to any reason whether obvious or not.

Wang et al (12) investigated the use of the induced membrane approach for reconstruction in 424 individuals who had been certified as "cured" of infection. In total, 52 patients suffered a recurrence of infection. Among these, 27.8% had the same bacterium as the initial infection.

In the current investigation, the majority of gram-negative organisms cultivated were methicillin-susceptible *S. aureus* (21%), with methicillin-resistant *S. aureus* accounting for 16% of samples. Methicillin-resistant *S. aureus* was the second most common causal organism of long bone infection, accounting for 16 of 81 positive culture growths, or roughly 20% of all bacterial cultures.

According to Garcia Pozo et al, tibia (62.1%) is the most usually affected bone, and *S. aureus* (58.5%) is the most prevalent pathogen.

In patients with osteomyelitis, treatment includes radical surgical debridement with or without antibiotic beads/cement spacers, as well as saucerization, sequestrectomy, and sinus tract excision.

The majority of studies advocate 6 weeks of intravenous (IV) antibiotic medication followed by further oral therapy for an unlimited period of time. However, reducing the length of IV antibiotic therapy to 1 week, followed by oral medication, did not result in increased remission rates (13).

*Staphylococcus aureus* is the most common organism responsible for both acute and chronic bone infections because it creates a biofilm that has the potential to rapidly develop antibiotic resistance and express virulence factors, regardless of the patient's immunological condition.

In our study, the average length of antibiotic administration was 6 weeks. Cloxacillin was the most commonly used antibiotic in 23 cases, Meropenem in 11, Septran in 8, Linezolid in 8, and Vancomycin in 7 cases.

Tice et al observed in their analysis of 454 cases of chronic osteomyelitis treated solely with outpatient-based parenteral therapy that both the initial pathogen and the type of antibiotic used for treatment determined recurrence. *Pseudomonas aeruginosa* presented a risk that was more than twice as high as that of *S. aureus* (14).

When necessary, a preoperative MRI may be performed to obtain a full view of bone and soft tissue, as well as to determine the precise extent of disease (15).

Jorge et al examined 192 patients suffering from posttraumatic osteomyelitis. They observed that recurrence was linked to elderly patients, intraoperative blood transfusions, and *P. aeruginosa* (16).

In our study, 25% of patients got a preoperative MRI, which was entirely at the discretion of the surgeon, taking into account the patient's financial situation, the extent and severity of the condition, and the surgical approach.

In our study, 19.3% of them had a recurrence of infection at any point in time over the 9<sup>th</sup> month follow-up, and 7.3% required repeat surgery during their hospital stay.

Many factors influenced the functional result following surgery, including mental state, number of previous treatments, degree of bone involvement, therapy decision for limb salvage, patient compliance to treatment, and so on.

In our study, we found a substantial clinical improvement in functional outcome in 7.6% of 91 patients who were followed up on 3 months after surgical debridement and 15.5% of 45 patients who were followed up on 6 months.

When compared to other groups, Cierny-Mader stage-1, hematogenous variety, and culture-negative osteomyelitis were characteristics that reduced the risks of recurrence.

Thirty-one patients were followed upto 9<sup>th</sup> month; 14 had chronic osteomyelitis, and 17 had infected non-union.

Patients with posttraumatic osteomyelitis had a greater recurrence rate (24.3%) than those with hematogenous (primary osteomyelitis; 10%).

Subramanyam et al investigated recurrence risk factors in 147 chronic osteomyelitis patients and found that polymicrobial growth and extended-spectrum beta-lactamase generating Enterobacteriaceae increased the likelihood of chronic osteomyelitis recurrence (17).

According to Petfield et al (18), recurrence is dictated by bone loss, the presence of devascularized bone, and the duration of antibiotic therapy. In a study of 142 cases with posttraumatic osteomyelitis, Chadayammuri et al (19) observed that polymicrobial development and a lack of skin cover were factors in poor prognosis. The administration of parenteral and oral antibiotics in sequence was associated with a lower frequency of ill outcomes.

According to Pozo et al, the length of chronic osteomyelitis for more than 3 months, bone exposure, and the need for further treatments (such as flaps) all contributed to the possibility of recurrence. The average time between debridement and the first signs of relapse was 10 months (2 weeks; 80 months).

The current study had short-term follow-up; therefore, definitive results about infection recurrence and long-term functional outcomes may not be conclusive.

The significance of this work is that it targets an infectious condition that requires early diagnosis at the community level because the burden of disease may interfere with the daily activities of people who belong to the lower-middle class. There is little data on relapse rates and functional outcomes following surgery in chronic osteomyelitis; therefore, this study can shed light on these facets of the disease.

## Conclusion

This study demonstrates a higher rate of infection recurrence or relapse in posttraumatic osteomyelitis. Chronic osteomyelitis relapsed at a rate of 19.8%. At 6 months, the functional outcome findings were better. Disease burden, financial issues, disability, and long duration of therapy all have an impact on the patient's mental state, which may further impact treatment adherence.

## Acknowledgement

None.

## Funding sources

The authors received no financial support for this article's research, authorship, or publication.

## Ethical statement

The study was performed per the Declaration of Helsinki, and written informed consent was obtained from the study participants. Institutional ethical approval (EC/23/01) was obtained for the study protocol.

## Conflicts of interest

The authors declare no conflict of interest in this research.

## Author contributions

DV and AT analyzed and interpreted the osteomyelitis patient's clinical and microbiological data. MS performed the surgical interventions and was a major contributor to writing the manuscript. KM supervised the total work and edited the manuscript. All authors read and approved the final manuscript.

## References

1. Uçkay I, Assal M, Legout L, Rohner P, Stern R, Lew D, et al. Recurrent osteomyelitis caused by infection with different bacterial strains without obvious source of reinfection. *J Clin Microbiol*. 2006;44(3):1194-6. [View at Publisher] [DOI] [PMID] [Google Scholar]
2. Wu Z, Zeng D, Yao J, Bian Y, Gu Y, Meng Z, et al. Research progress on diagnosis and treatment of chronic osteomyelitis. *J Chin Med Sci*. 2019;34(3):211-20. [View at Publisher] [DOI] [PMID] [Google Scholar]

3. Masrouha KZ, Raad ME, Saghih SS. A novel treatment approach to infected non-union: of long bones without systemic antibiotics. *Strategies Trauma Limb Reconstr.* 2018;13(1):13-8. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
4. Lowenberg DW, DeBaun M, Suh GA. Newer perspectives in the treatment of chronic osteomyelitis: A preliminary outcome report. *Injury.* 2019;50(1):S56-61. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
5. Geraghty T, Laporta G. Current health and economic burden of chronic diabetic osteomyelitis. *Expert Rev Pharmacoecon Outcomes Res.* 2019;19(3):279-86. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
6. Ellur V, Kumar G, Sampath JS. Treatment of chronic hematogenous osteomyelitis in children with antibiotic impregnated calcium sulphate. *J Pediatr Orthop.* 2021;41(2):127-31. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
7. Bose D, Kugan R, Stubbs D, McNally M. Management of infected non-union: of the long bones by a multidisciplinary team. *Bone Joint J.* 2015;97(6):814-7. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
8. Biruk WL, Wubshet K. Chronic Osteomyelitis at Tikur Anbessa Hospital, Addis Ababa University, Ethiopia. *East Cent Afr J.* 2007;12:33-41. [[View at Publisher](#)] [[Google Scholar](#)]
9. Chrousos GP. Stress, chronic inflammation, and emotional and physical well-being: concurrent effects and chronic sequelae. *J Allergy Clin Immunol.* 2000;106(5):S275-91. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
10. Uçkay I, Jugun K, Gamulin A, Wagener J, Hoffmeyer P, Lew D. Chronic osteomyelitis. *Curr Infect Dis Rep.* 2012;14:566-75. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
11. Garcia del Pozo E, Collazos J, Carton JA, Camporro D, Asensi V. Factors predictive of relapse in adult bacterial osteomyelitis of long bones. *BMC Infect Dis.* 2018;18(1):635. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
12. Wang X, Wang S, Fu J, Sun D, Shen J, Xie Z. Risk factors associated with recurrence of extremity osteomyelitis treated with the induced membrane technique. *Injury.* 2020;51(2):307-11. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
13. Okumura N, Hayakawa K, Yamamoto K, Yamada G, Mezaki K, Ohmagari N. Effectiveness of oral cephalexin in antibiotic-course completion for methicillin-susceptible *Staphylococcus aureus*-induced bacteremic vertebral osteomyelitis. *BMC Infect Dis.* 2023;23(1):307. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
14. Tice AD, Hoaglund PA, Shoultz DA. Risk factors and treatment outcomes in osteomyelitis. *J Antimicrob Chemother.* 2003;51(5):1261-8. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
15. Marais LC, Ferreira N, Aldous C, Le Roux TL. The management of chronic osteomyelitis: Part I-Diagnostic work-up and surgical principles. *SA Orthopaedic Journal.* 2014;13(2):42-8. [[View at Publisher](#)] [[Google Scholar](#)]
16. Jorge LS, Chueire AG, Fucuta PS, Machado MN, Oliveira MG, Nakazone MA, et al. Predisposing factors for recurrence of chronic posttraumatic osteomyelitis: a retrospective observational cohort study from a tertiary referral center in Brazil. *Patient Saf Surg.* 2017;11:17. . [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
17. Subramanyam KN, Mundargi AV, Prabhu MV, Gopakumar KU, Gowda DA, Reddy DR. Surgical management of chronic osteomyelitis: Organisms, recurrence and treatment outcome. *Chinese. Chin J Traumatol.* 2023;26(4):228-35. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
18. Petfield JL, Tribble DR, Potter BK, Lewandowski LR, Weintrob AC, Krauss M, et al. Is bone loss or devascularization associated with recurrence of osteomyelitis in wartime open tibia fractures?. *Clin Orthop Relat Res.* 2019;477(4):789-801. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]
19. Chadayammuri V, Herbert B, Hao J, Mavrogenis A, Quispe JC, Kim JW, et al. Factors associated with adverse postoperative outcomes in patients with long bone post-traumatic osteomyelitis. *Eur J Orthop Surg Traumatol.* 2017;27(7):877-82. [[View at Publisher](#)] [[DOI](#)] [[PMID](#)] [[Google Scholar](#)]

### How to Cite:

Shaik MB, Mallam KK, Vurundhur D, Talari AB. A study of relapse rate and functional outcome after surgical debridement in patients with chronic osteomyelitis. *JCBR.* 2023;7(4):10-4.



© The author(s)