

## Case Report

### Chronic Diarrhea as the Sole Symptom of HIV Infection

Mehdi Alimadadi<sup>1\*</sup>, Sina Mohajernoeei<sup>2</sup>, Hanieh Raghimi<sup>2</sup>,

1. Golestan Research Center of Gastroenterology and Hepatology, Golestan University of Medical Sciences, Gorgan, Iran

2. Department of Medical Sciences, Faculty of Medicine, Golestan University of Medical Sciences, Gorgan, Iran

\***Correspondence:** Dr Mehdi Alimadadi, Golestan Research Center of Gastroenterology and Hepatology, Golestan University of Medical Sciences, Gorgan, Iran.

Email: alimadadi1360@yahoo.com

Tel: +989126858962

Orchid id: 0000-0002-2102-7320

**Received** August 4, 2021

**Accepted** September 14, 2021

#### ABSTRACT

**Background:** Acquired immunodeficiency syndrome (AIDS) is a disease caused by the human immunodeficiency virus (HIV) that can affect all systems of the human body. Considering that there is no definitive treatment for AIDS, timely and accurate diagnosis is important. Symptoms of HIV infection may sometimes be nonspecific and should always be considered in high-risk individuals. We herein report a patient with HIV who was presented with chronic diarrhea as the sole symptom of HIV Infection.

**Case description:** A 40 years old man was admitted to hospital with complain of diarrhea that had started seven months ago. He also experienced extensive weight loss and complained of vomiting and abdominal pain in the past month without any fever. Abdominal computed tomography scan and colonoscopy and stool examination were normal. However, biopsy indicated active colitis. Laboratory tests for HIV, functional expression of the lymphoid chemokines, and anti-tissue transglutaminase antibody were negative, but hepatitis C viral load was positive. Because the patient had many tattoos on his body that were done in unsanitary places, AIDS was suspected. The related test was requested, and the result was positive. Finally the patient was prescribed with co-trimoxazole and referred to an AIDS consultation center.

**Conclusion:** We presented a case of HIV infection with chronic diarrhea as the only symptom without any others clinical findings. Our observations suggest that for better diagnosis and management of HIV infection, it is important to consider chronic diarrhea as one of the important complications of HIV even when not accompanied with any other symptom.

**Keywords:** Acquired immunodeficiency syndrome, diarrhea, HIV, colon

**DOI:** 10.29252/Jcbr.5.3.21



This work is licensed under a Creative Commons Attribution 4.0 License.

© The authors

## INTRODUCTION

Acquired immunodeficiency syndrome (AIDS) is a serious disease that can affect all systems of the human body. The disease is mainly transmitted through bodily fluids and unprotected sexual intercourse. Symptoms of human immunodeficiency virus (HIV) infection depend on the phase of infection; therefore, HIV infected individuals can be asymptomatic or present few clinical manifestations for years before progression to AIDS. This disease weakens the immune system, making the body susceptible to other infections and treatment more challenging, leading to increased risk of certain comorbidities and mortalities in HIV-infected individuals. In addition, AIDS can involve different systems of the body including skin, nervous system, lymph glands, lungs, mucous membrane, and gastrointestinal (GI) tract (1-4). One of the common GI symptoms is diarrhea, particularly secretory diarrhea that can occur in all stages of HIV infection (5). This type of diarrhea can cause several complications and lead to death if left untreated (5-7).

AIDS can be diagnosed through blood or saliva testing, and CD4 cell count and viral load (HIV RNA) are useful for determining the stage of the disease (8). Stool and endoscopic examinations play main roles for differential diagnosis and evaluation of diarrhea in HIV infection (9).

Currently, there's no definitive cure for AIDS and available medications are for controlling the course of HIV infection and preventing complications (10).

Considering that incidence of diarrhea without other complications is rare and dangerous in HIV patients, in this study, we describe a case of AIDS who was presented with diarrhea only.

## CASE PRESENTATION

A 40-year-old male patient admitted to hospital with complaints of chronic diarrhea, which had begun seven months ago. At first, he had suffered two episodes of loose stool per day, which then increased to four. The patient's stool was yellow and in liquid


form, with no trace of blood. He also reported a weight loss of about 30 kg over the last six months. He complained of vomiting and abdominal pain in the last month as well but had no complaints of fever.

The patient had been evaluated in another hospital for diarrhea three months ago, and some areas of colon were erythematic with abnormal vascular patterns in the colonoscopy examination. He had then undergone biopsy examination through which a mild active colitis was found, and the patient was recommended for follow-up evaluations.

In immunological assessments, HIV (types 1 and 2), functional expression of the lymphoid chemokines and anti-tissue transglutaminase antibody were reported negative but we considered the possibility of false negative results. In addition, anti-hepatitis C virus (HCV) antibody was reported in a borderline state. Laboratory findings indicated presence of HCV viral load. The patient was then treated using a regimen of ciprofloxacin and metronidazole, but symptoms remained unaffected.

In physical examination, abdomen was soft and without tenderness, and bowel sounds were increased. Vital signs were normal. Colonoscopy examination did not show any gross abnormalities. A biopsy for microscopic colitis was performed, which revealed active colitis, without evidence of crypt destruction or lymphocytic/collagenous colitis (Figure 1). Spiral abdominopelvic computed tomography (CT) scan with intravenous (IV) contrast and abdominal sonography revealed no abnormalities. Stool examination and stool culture were normal.

Considering the fact that all paraclinical measures were normal and the patient had a history of tattooing his body in unsanitary places, HIV testing using PCR was performed on the patient which was reported positive. Ultimately the patient was prescribed with co-trimoxazole and referred to an AIDS consultation center.



**Figure1. Hematoxylin and eosin staining of colon biopsy.** Besides normal transverse colon tissue, histologic microscopy shows round straight crypts and inflamed lamina propria involved by acute and chronic inflammation. There are cryptitis and crypt abscess formation. The images were taken under  $\times 40$  (a),  $\times 10$  (b), and  $\times 4$  (c) magnifications.

## DISCUSSION

Opportunistic infections, uncommon malignant neoplasm, and persistent immunologic deficiency are AIDS allmarks (11). The gastrointestinal problems associated with AIDS include diarrhea, nausea, vomiting, bloating, abdominal discomfort, and changes in body weight. The nature of AIDS-associated diarrhea can be non-infectious and infectious. Bacteria, fungi, viruses, and protozoa are opportunistic pathogens that can cause infectious diarrhea. Since the advent of highly-active antiretroviral therapy, the incidence of infectious and non-infectious diarrhea has decreased and increased, respectively.

Exposure of naïve B and T cells to antigens in the gut-associated lymphoid tissue leads to their maturation into plasma cells and memory T cells. This process generates a baseline inflammatory state to provide chemokines and adhesion molecules, thereby attracting lymphocytes towards mucous layer. As HIV multiply inside lymphocytes, the GI tract becomes one of the most targeted areas and HIV-associated diarrhea can occur (12).

Presence of diarrhea as the only complication of HIV makes the disease difficult to diagnose. Given the importance of this symptom, prompt and timely diagnosis is essential.

## CONCLUSION

Altogether our observations show that chronic diarrhea could be considered as the sole clinical manifestation of HIV-infected patients.

## ACKNOWLEDGMENTS

None.

## DECLARATIONS

### Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

### Ethics approvals and consent to participate

A written informed consent was obtained from the patient. The patient received all the necessary clinical treatments and the study involved no intervention.

### Conflict of interest

The author declares that there is no conflict of interest.

## REFERENCES

1. Price WA, Forejt J. Neuropsychiatric aspects of AIDS: a case report. *General hospital psychiatry*. 1986 Jan 1;8(1):7-10. [[View at Publisher](#)] [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
2. Gottlieb MS, Schroff R, Schanker HM, Weisman JD, Fan PT, Wolf RA,

Saxon A. Pneumocystis carinii pneumonia and mucosal candidiasis in previously healthy homosexual men: evidence of a new acquired cellular immunodeficiency. *New England Journal of Medicine*. 1981 Dec 10;305(24):1425-31. [[View at Publisher](#)] [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

3. HIV.gov. What are HIV and AIDS?. June 05, 2020 .<https://www.hiv.gov/hiv-basics/overview/about-hiv-and-aids/what-are-hiv-and-aids>.

4. Scutchfield FD, Benenson AS. AIDS update. *Postgrad Med*. 1989 Mar;85(4):289-94. [[View at Publisher](#)] [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

5. Patel TS, Crutchley RD, Tucker AM, Cottreau J, Garey KW. Crofelemer for the treatment of chronic diarrhea in patients living with HIV/AIDS. *HIV/AIDS (Auckland, NZ)*. 2013;5:153. [[View at Publisher](#)] [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

6. sTwisha S Patel, Rustin D Crutchley, Anne M Tucker, Jessica Cottreau, Kevin W Garey. Crofelemer for the treatment of chronic diarrhea in patients living with HIV/AIDS. *HIV AIDS (Auckl)*. 2013; 5: 153-162. [[View at Publisher](#)] [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

7. Barrett KE, Keely SJ. Chloride secretion by the intestinal epithelium: molecular basis and regulatory aspects.

*Annual review of physiology*. 2000 Mar;62(1):535-72. [[View at Publisher](#)] [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

8. Hardy WD. *Fundamentals of HIV Medicine* 2019. Oxford University Press, USA; 2019 Jun 10. [[View at Publisher](#)] [[Google Scholar](#)]

9. Noor A, Bedi D, Yassin MH. Multiorganism HIV-associated diarrhea and telescoping bowel. In *Baylor University Medical Center Proceedings* 2020 Jul 2 (Vol. 33, No. 3, pp. 386-388). Taylor & Francis. [[View at Publisher](#)] [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

10. Pollack TM, et al. Primary care of the HIV-infected adult. <https://www.uptodate.com/contents/search>. Accessed Dec. 6, 2019

11. Sánchez-Mejorada G, Ponce-de-Leon S. Clinical patterns of diarrhea in AIDS: etiology and prognosis. *Revista de investigacion clinica; organo del Hospital de Enfermedades de la Nutricion*. 1994 May 1;46(3):187-96. [[View at Publisher](#)] [[PubMed](#)] [[Google Scholar](#)]

12. Hall VP. Common Gastrointestinal Complications Associated with Human Immunodeficiency Virus/AIDS: An Overview. *Critical care nursing clinics of North America*. 2017 Dec 6;30(1):101-7. [[View at Publisher](#)] [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

**How to Cite:** alimadadi M, Mohajernei S, Raghimi H. Chronic Diarrhea as the Sole Symptom of HIV Infection: A Case Report. *Journal of Clinical and Basic Research*. 2021; 5 (3) :21-24