

***In vivo* and *in silico* investigation of effects of ethanol extract of *Moringa oleifera* leaves on female fertility, using fruit flies and molecular docking**

Ben Enoluomen Ehigiator¹ , Uchechukwu Loveth Iyanyi^{1*} , Kelechi Samuel Mobisson² , Onuchkwu Imelda Ukata¹ 

1. Department of Pharmacology and Toxicology, Faculty of Pharmacy, Madonna University, Elele, Rivers State, Nigeria

2. Department of Human Physiology, Faculty of Basic Medical Sciences, Madonna University, Elele, Rivers State, Nigeria

* Correspondence: Uchechukwu Loveth Iyanyi. Department of Pharmacology and Toxicology, Faculty of Pharmacy, Madonna University, Elele, Rivers State, Nigeria. Tel: +2348135008001, Email: uchechukwuiyanyi@yahoo.com

Abstract

Background: Infertility is an important issue for couples that may cause various psychological and emotional problems. Female infertility disorders play a major role in approximately 50-80% of the causes of infertility in various areas in Nigeria. *Moringa oleifera* has been proposed as a plant with female fertility enhancement effects. The objective of this study was to assess the fertility-improving effects of ethanol extract of *M. oleifera* leaf and to determine the phytochemical components causing these effects by *in silico* analyses.

Methods: The *in vitro* effects on fertility were evaluated using *Drosophila melanogaster* (fruit fly) because of its genetic similarities to humans. The copulation duration, mating latency, and the number of emergences from the fruit fly after mating were determined. Three doses (0.025%, 0.05%, and 0.1% w/w) of the *M. oleifera* ethanol extract were administered to three different groups, while a control group only received feed mixed with ethanol. For *in silico* studies, 62 compounds were obtained from the PubChem library by mining compounds from articles related to *M. oleifera*. Next, a ligand library was generated and docked against various targets of interest (estrogen, progesterone, kisspeptin, liver X, PPAR γ , and 15-PGDH receptors as well as 17 β hydroxysteroid dehydrogenase and insulin-degrading enzymes) which have female fertility-enhancing effects.

Results: The *in vivo* experiments showed that *M. oleifera* had no effect on copulation duration and mating latency, but interestingly, it enhanced the fertility/emergence of the treated fruit flies. *In silico* studies suggested that phytochemicals such as rutin, marumosioid B, myricetin, and quercetin showed docking scores that may well support previous works on *M. oleifera* enhancement of female fertility.

Conclusion: The results showed that *M. oleifera* can enhance fertility in female fruit flies.

Article History

Received: 25 April 2023

Received in revised form: 22 August 2023

Accepted: 22 August 2023

Published online: 24 November 2023

DOI: [10.29252/JCBR.7.3.1](https://doi.org/10.29252/JCBR.7.3.1).

Keywords

[Moringa oleifera](#)
[Fertility agents, female](#)
[Drosophila melanogaster](#)
[3 \(or 17\)-beta-hydroxysteroid dehydrogenase](#)
[In silico model](#)

Article Type: Original Article



Highlights

What is current knowledge?

The high level of fertility in treated flies was dose-dependent.

What is new here?

Rutin and myricetin present in *M. oleifera* are responsible for the female fertility enhancement effect.

Introduction

Reproduction is one of the most important traits of all living organisms, which is crucial for the continuation of life. Infertility has continued to be an issue of concern for couples and sexually active people who want to have children, and their search for a therapeutic solution has proven to be sometimes fruitless. A condition of the reproductive system known as infertility is characterized by the inability to achieve a clinical pregnancy after at least a year of regular, unprotected sex without complications like lactation or postpartum amenorrhea. Infertility affects about 10% of people of reproductive age (1) and 15% of couples globally, 50% of which are female-related alone (2). This condition can be related to both innate factors (genetic, immunological, anatomical, and hormonal disorders) and external factors (obesity, infections after childbirth or surgery, tuberculosis of the pelvis, and obesity) (3). For couples, infertility is a major cause of concern, and in various areas of Nigeria, female infertility accounts for 50-80% of all divorces (4). Female infertility has been linked to several medical conditions including pelvic inflammatory disease, polycystic ovarian syndrome, endometriosis, premature ovarian failure, uterine fibroids, and environmental toxins such as glues, volatile organic solvents, silicones, organic dust, pesticides, and physical agents (5). Ovulation and uterine problems, tubal blockages, hormonal imbalances, prior tubal ligations, and weight changes that can lead to ovarian dysfunction are possible additional causes of infertility in females (6). In addition, a low body fat percentage impairs the production of estrogen, which results in an aberrant ovulation cycle and irregular menstruation (7).

Moringa oleifera is a plant from the family Moringaceae with a wide range

of medicinal applications (8). It has been also used in the production of pharmaceutical and industrial products and therefore has been regarded as the "miracle tree" (9, 10). The phytochemical screening of *M. oleifera* leaf extract has demonstrated the presence of alkaloids, saponins, tannins, steroids, phenolic acids, glucosinolates, flavonoids, and terpenes (11). Numerous researches in Nigeria have shown that *M. oleifera* contains both macro- and micronutrients that can be used for the treatment of female infertility (12-14). The plant has been thought to improve female fertility by increasing the secretion or availability of ovarian hormones, such as progesterone and estrogen, which are in charge of pregnancy maintenance, ovulation, implantation, and delivery (15, 16). This study aimed to assess the impact of ethanol extract of *M. oleifera* leaf on the reproductive potential of fruit flies. Several organisms share the fruit fly's basic genetic mechanism, and hundreds of closely similar species have been thoroughly investigated for decades, with a wealth of literature available (17). In this study, molecular docking analysis was used to identify *M. oleifera*'s molecular target activity.

Methods

The materials used include breeding and treatment vials, a micropipette, empendox tubes, cotton wool, a stirring rod, a spatula, a pot, ethanol, sample bottles, a beaker, a measuring cylinder, distilled water, filter paper, funnel, an electric cooker, water bath, and blender.

Preparation of *M. oleifera* leaf

Fresh *M. oleifera* leaves were purchased from a nearby plantation in Mararaba, Nasarawa State, Nigeria. Botanists identified the plant leaves, assigned them to herbarium number 08, and let them air dry at room temperature. The dried leaves were powdered and soaked in 250 ml of ethanol for 72 hours. After filtration, the mixture was dried in a water bath (B-Bran Scientific and Instrument Company, England) at 60°C. As described previously (18), the brownish residue was cleaned and refrigerated until used in an airtight bottle.

Feed preparation and dose concentrations

Feed preparation was carried out as explained previously (19). The feed was then well stirred into a mixture, weighed into the treatment vials, covered, and left to cool naturally.

Preparation of offspring used for the study

Flies were bred on corn agar media. The eggs hatched into adult fruit flies after 7 days of breeding. After approximately 10 days of breeding, the stock generation of flies was removed, and the daughter generation (offspring) was allowed to acclimatize at $25 \pm 2^\circ\text{C}$ until the needed number of flies for treatment was obtained. Four groups of 25 flies (both sexes) were used for the investigation. Each group received three duplicates and received the ethanolic extract of *M. oleifera* at three distinct concentrations: 0.025%, 0.05%, and 0.1% w/w. Dead flies were counted and recorded every day, and the percentage of surviving flies was calculated up until the point at which all insects were dead.

Survival test and lethal concentration 50 (LC₅₀) determination

Four separate groups (A, B, C, and D), each containing 25 flies, received 0, 0.05%, 0.1%, and 0.2% w/w of the extract incorporated in the feed for 28 days. The groups were examined daily for dead flies. Next, the subacute median lethal concentration (LC₅₀) was determined as described in a previous study (19).

Fruit fly preparation

The fruit flies for the fertility and survival studies of *M. oleifera* consisting of both males and females were obtained from the Pharmacology and Toxicology Department of Madonna University, Elele, Rivers State. The fruit flies were bred. The study was carried out after obtaining the approval of the Animal Research Ethics Committee, Madonna University, Elele, Nigeria (EC/FP/023/14).

Mating experiment

Less than 8 hours after enclosure, 25 male flies were randomly divided into 4 distinct vials (n=6) marked A (untreated), B, C, and D. Various concentrations of the ethanolic root extract of *M. oleifera* (0.025%, 0.05%, and 0.1% w/w) were added to the feed for groups B-D. One virgin female and one treated virgin male were coupled in one vial for each group. After the flies were coupled, the mating activity was observed for one hour. For each of the 24 pairs, the length of the copulation was scored.

Fertility studies

The female flies were allowed to lay eggs for 2 days after each pair was moved to a fresh agar-corn meal diet. Then, the pair was moved into another treatment vial with fresh feed for an additional 2 days. The parent flies were then disposed of by placing them in a methanol-filled bottle. After 8 days, the offspring from each replicate vial labeled A1–A6, B1–B6, C1–C6, and D1–D6 were scored for 7 days until no more emergence occurred (19).

Ligand library generation

To create the ligand library, identified secondary metabolites of *M. oleifera* were selected from published literature. In Standard Database Format (2D), 60 secondary metabolites were obtained from the NCBI PubChem database. The resulting ligand library was produced using the docking software Maestro (Schrodinger suite, version 2018-1b) as described in a previous study (20).

Protein preparation

With the help of their respective protein databank (PDB) IDs, a variety of targets were obtained from the PDB along with their associated co-ligands. These included the estrogen receptor co-ligand (1FDT), progesterone receptor co-ligand (1A28), liver X receptor co-ligand (3LOE), peroxisome proliferator-activated receptor gamma (PPARG) co-ligand (6KOT), the standard agent for 15-hydroxyprostaglandin dehydrogenase (15-PGDH) SW 033291 (2GDZ), 17 beta-hydroxysteroid dehydrogenase co-ligand (5L7Y), and insulin-degrading enzyme co-ligand amylin (3HGZ) as described previously (21) using the Protein Preparation Wizard (22).

Statistical analysis

GraphPad Prism (version 5.01) was used for the statistical analysis.

Results

Survival studies of *M. oleifera* on *D. melanogaster*

The results revealed that the survival rates of the *M. oleifera* ethanol leaf extract treatment groups (B, C, and D) did not differ significantly compared to that of the control group A (Figure 1).




Figure 1. Survival studies of *M. oleifera* on *D. melanogaster*

Effect of *M. oleifera* on mating latency

Mating latencies of the treated groups differed significantly compared with the control group. However, the mating latencies of the treated group were longer at the lowest dose (50 mg/kg, group B) (Figure 2).




Figure 2. Effect of *M. oleifera* on mating latency

Copulation duration

There was a great decline in the copulation durations in a dose-dependent manner. No mating or copulation was observed within an hour of observation in group D (the highest-dose treatment group) (Figure 3).




Figure 3. Effect of *M. oleifera* on copulation duration (* $P < 0.05$ *M. oleifera* versus the control group)

Effect of *M. oleifera* on the emergence of flies

The result showed a paradoxical but interesting effect. We detected a high fertility level and fecundity level in the treated flies. Nevertheless, there was no statistically significant difference between the groups in terms of fertility level (Figure 4).




Figure 4. Effect of *M. oleifera* on the emergence of flies

Molecular docking study

Figures 5 to 12 show the results of molecular docking study on estrogen receptor, progesterone receptor, kisspeptin receptor, PPARG, 15-PGDH, 17β hydroxysteroid dehydrogenase, insulin-degrading enzyme, and liver X receptor.




Figure 5. Heat map representation of docking result for compound interaction with the estrogen receptor (-4 kcal/mol) to red (-12 kcal/mol)




Figure 6. Heat map representation of docking result for compound interaction with the progesterone receptor (-4kcal/mol) to red (-12kcal/mol)




Figure 7. Heat map representation of docking result for compound interaction with the kisspeptin receptor (-0kcal/mol) to red (-9kcal/mol)




Figure 8. Heat map presentation of docking result for compound interaction with PPARγ (-4kcal/mol) to red (-13kcal/mol)




Figure 9. Heat map presentation of docking resulting from compound interaction with 15-PGDH (-4kcal/mol) to red (-12kcal/mol)




Figure 10. Heat map presentation of docking resulting from compound interaction with 17β hydroxysteroid dehydrogenase (-4kcal/mol) to red (-12kcal/mol)




Figure 11. Heat map presentation of docking resulting from compound interaction with the insulin-degrading enzyme (-4kcal/mol) to red (-10kcal/mol)




Figure 12. Heat map presentation of docking resulting from compound interaction with Liver X receptor (-4kcal/mol) to red (-13kcal/mol)

Discussion

For couples and sexually active people who desire to have children, infertility has continued to be a major hurdle and health issue. In underdeveloped nations where manufactured treatments are expensive and difficult to obtain, the use of medicinal plants for infertility treatment might be a suitable alternative (23). Orthodox drugs such as clomiphene, tamoxifen, gonadotropins, metformin, letrozole, and bromocriptine have some undesirable effects. These side effects have prompted research on the use of medicinal plants to solve female infertility problems. Numerous investigations on the effect of *M. oleifera* on female fertility have revealed that the plant contains phytoestrogens, which exert estrogen-like effects. Treatment with *M. oleifera* ethanol extract on rats for 30 days resulted in a significant rise in luteinizing hormone and estrogen contrary to the follicle-stimulating hormone level, which remains unchanged (15). The extract's activation of estrogen receptors is assumed to be the cause of the elevated levels of estrogen and luteinizing hormone. The compounds present in the extract can bind to estrogen receptors to bring about estrogenic responses (15).

According to the literature, the effects of ethanol extract of *M. oleifera* on the ovary may be secondary to their basic actions on the hypothalamus or pituitary gland. This results in the amplification of normal gonadotropin release (16, 24). The administration of *Moringa* leaves at 200 and 500 mg/kg body weight for 30 days significantly increased the number of implantation sites on the 8th day of pregnancy in rats. No implantation site was observed in the high- and low-dose group of rats administered with other parts of the *Moringa*, namely the seed, flower, root, and seed. The findings suggested that the consumption of *M. oleifera* leaf extract increases female reproductive function and fecundity in rats, as noticed by the increase in the number of embryos successfully implanted in the rats' uterus. According to the results of this study, *Moringa* leaves may improve female reproductive function by boosting the availability or release of ovarian hormones such as progesterone and estrogen. This suggests that *M. oleifera* ingestion is safe and does not result in fetal abortion.

In the present study, fruit fly was used for in vivo experimentations because of their genetic similarities to humans, very short life cycle, and production of a large number of offspring, which allow sufficient data collection. We determined the survival of the fruit fly using different concentrations of the extract and measured copulation duration, mating latency, and emergence of the flies after mating. The lack of significant difference in the survival rate of subjects between the treated groups and the control group might indicate that *M. oleifera* does not affect the survival rate of fruit flies. The mating latencies of the treated groups were longer; therefore, its lowest dose may be safer. This may be an indication that this agent is not likely to have aphrodisiac properties due to a notable decline in the copulation durations in a dose-dependent manner. The emergence of fruit flies could indicate the fertility enhancement effect, showing that there was a high level of fertility and fecundity in treated flies in a dose-dependent pattern. These positive results provided the need to ascertain the different targets to predict the pro-fertility potentials of *M. oleifera* component by conducting in silico studies. As mentioned previously, *M. oleifera* has been adjudged to contain macro and micronutrients that are potentially beneficial for the treatment of female infertility (12-14). Phytochemical screening of *M. oleifera* has previously revealed the presence of saponins, alkaloids, steroids, tannins, steroids, phenolic acids, glucosinates, terpenes, and flavonoids (11). In this study, virtual screening of *M. oleifera* constituents from the literature was carried out, and the following compounds were obtained: rutin, apigenin, isorhamnetin, quercetin, luteolin, genistein, daidzein, myricetin, epicatechin, sinalbin, salicylic acid, gentistic acid, syringic acid, caffeic acid, sinapic acid, o-coumaric acid, p-coumaric acid (25), isoquercetin, astragalol, cryptochlorogenic acid (26), procyanidins (27), beta-amyrin, alpha-amyrin, neochlorogenic, rhamnetin (28), vicenin-2 (29), benzyl isothiocyanate, oleic acid, myristic acid, palmitic acid, isothiocyanate (30), E-zeaxanthin, Z-lutein, E-lutein, E-luteoxanthin, beta-carotene (31), gallic acid, ellagic acid, ferulic acid, chlorogenic acid (32), paullinic acid, behenic acid, linoleic acid, beta-sitosterol, arachidic acid, stearic acid, palmitoleic acid, stigmaterol, campesterol (33), lupeol, vanillin (34), eugenol (35), niazirin, niaziridin (36), rhamnopyranoside (37), niazimicin (38), and niaziminin (39). Some phytochemical compounds present in *M. oleifera*, such as rutin, quercetin, isorhamnetin, lupeol, vicenin-2, chlorogenic acid, vanillin, isoquercetin, and cryptochlorogenic acid have been shown to possess fertility effects by enhancing ovarian folliculogenesis to bring about ovulation, enhance steroidogenesis by binding to their nuclear receptors, regulating some enzymes of steroidogenesis or regulating the activities of key enzymes involved in their metabolism (40). Such regulator effect on the reproductive function can be directly linked to its influence on the hypothalamic-pituitary-ovarian axis by stimulating or inhibiting the enzymes and receptors involved in this process and steroidogenesis disrupting the hormonal functioning of the hypothalamus and pituitary gland. Some receptors and enzymes must be stimulated or inhibited to improve female fertility since they are responsible for follicle development, oocyte maturation, fertilization rates, pregnancy outcome, embryo development, and early implantation. The receptors with this fertility effect include progesterone receptor, estrogen receptor, kisspeptin receptor, liver X receptor, and peroxisome proliferator activators receptor, and the enzymes include 15-hydroxy prostaglandin dehydrogenase, 15-beta hydroxysteroid dehydrogenase, and insulin-degrading enzyme.

The fertility potential of rutin, cryptochlorogenic acid, luteolin, myricetin, Isorhamnetin, quercetin, marumosi B, vicenin-2, isoquercetin, sinalbin, and astragalol present in *M. oleifera* was established by the stimulatory effect on the progesterone, estrogen, Kisspeptin, liver X, and PPARG receptors and the inhibitory effects on 15-hydroxy prostaglandin dehydrogenase, 15-beta hydroxysteroid dehydrogenase, and insulin-degrading enzymes, all of which are involved in female fertility. Molecular docking studies are based on the fact that compounds possess the capacity to bind to amino acid residues, which is a function of protein conformation and the assumed pose of the ligand (41). The library of phytochemicals derived was docked against the five target proteins, which may be remedial in infertility in females.

Estrogen and progesterone receptors are essential since both are the functional key hormones for female fertility. In other words, stimulation of these hormones enhances fertility in females. In the docking studies, heat map representation revealed that myricetin has a high stimulatory docking affinity

for estrogen receptors significantly higher than that for the co-crystallized ligand at -12.155kcal/mol. With a docking score of -12.362 kcal/mol, epicatechin showed a similar potential to myricetin to that of the co-ligand, which was -11.933 kcal/mol. Quercetin may also stimulate progesterone receptors as it possesses a high docking score of -12.477 kcal/mol, and myricetin at -12.086kcal/mol is more than the co-ligand at -11.988 kcal/mol. When stimulated, the kisspeptin receptor also causes the secretion of gonadotropins, including luteinizing hormone (LH) and follicle-stimulating hormone, which enhances female fertility (42). The docking studies revealed that rutin stimulates the kisspeptin receptor more than the co-ligand at 11 kcal/mol.

Peroxisome proliferator-activated receptors also play a role in female fertility and reproduction, which increases the pituitary production of prolactin and LH (43), thereby assisting folliculogenesis. The heat map representation revealed that vicenin-2 may stimulate PPARG significantly and even more than the co-crystallized ligand at -13.155kcal/mol, while rutin may stimulate PPARG at -11.008 kcal/mol compared to the co-ligand at -9.39 kcal/mol.

Stimulation of prostaglandin E2 receptor plays a role in female fertility by increasing LH levels, thereby enhancing ovulation, implantation, and menstruation (44). Enzymes that degrade prostaglandins such as 17 β hydroxysteroid dehydrogenase and 15-PGDH also play a role in enhancing female fertility. Inhibiting this enzyme that metabolizes prostaglandin leads to an increased level of prostaglandin and thus can facilitate the actions of prostaglandin in fertility enhancement in females. Heat map representation also revealed that rutin may inhibit the 15-PGDH enzyme significantly even more than the co-crystallized ligand at -12.428 kcal/mol, and procyanidins had a close effect on rutin with inhibition of 15-PGDH at -12.428 kcal/mol compared to the co-ligand at -9.635 kcal/mol. Rutin also may inhibit 17 β hydroxysteroid dehydrogenase at -12.813 kcal/mol and procyanidins at -11.497 kcal/mol more than the co-ligand at -7.989 kcal/mol.

Insulin-like peptide 3 (INSL3) receptor plays a role in increasing progesterone production (45) and oocyte maturation (46). Insulin-degrading enzymes must be inhibited to increase the level of insulin for this effect. The results obtained revealed that rutin may inhibit insulin-degrading enzymes significantly even more than the co-crystallized ligand at -10.134 kcal/mol. Marumosi B may also inhibit insulin-degrading enzymes at -9.111 kcal/mol compared to the co-ligand at -8.236 kcal/mol.

Stimulation of the liver X receptor influences cholesterol metabolism, which is the precursor for ovarian steroid biosynthesis (47). The results showed that isoquercetin stimulates the liver X receptor at -13.813 kcal/mol, marumosi B at -13.016 kcal/mol, and astragalol at 3.129 kcal/mol even more than the co-ligand at -11.962 kcal/mol.

Our results obtained from the molecular docking studies indicate that rutin and myricetin found in *M. oleifera* have the potential to produce the best effect in almost all receptors and enzymes involved in the fertility process.

Conclusion

The ethanol extract of *M. oleifera* leaf shows less effect in copulation and mating latency but affects the number of fruit flies' emergence after mating, supporting the claim that *M. oleifera* has a female fertility-enhancing effect. Effective stimulation of the estrogen, progesterone, and PPARG receptors and inhibition of 15-PDGH, insulin-degrading, and 17 β hydroxysteroid dehydrogenase enzymes indicate the potential of this extract for female fertility enhancement, which can be mainly attributed to rutin and myricetin.

Acknowledgement

The researchers at the Centre for Bio-computing and Drug Development at Adekunle Ajasin University in Ondo and the Pharmacology and Pharmacognosy Department of Madonna University in Nigeria are both gratefully acknowledged by the authors.

Funding sources

The study received no funding from the government, business, or non-profit sectors.

Ethical statement

The study was carried out after obtaining the approval of the Animal Research Ethics Committee, Madonna University, Elele, Nigeria (EC/FP/023/14).

Conflicts of interest

There is no conflict of interest regarding the publication of this article.

Author contributions

E B.E designed and statistically analyzed the data for the study. The initial draft

of the manuscript was written by S. K. M. Laboratory tests and literature searches for methodology were managed by O.U.I. and U.L.I. The final manuscript was reviewed and approved by all writers.

References

- Zegers-Hochschid F, Adamson GD, De Mouzon J, Ishihara D, Mansour R, Nygren K, et al. The International Committee for Monitoring Assisted Reproductive Technology (ICMART) and the World Health Organization (WHO) revised glossary on ART terminology, 2009. *Hum Reprod.* 2009;24(11):2683-7. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Agarwal A, Mulgund A, Hamada A, Chyatte MR. A unique view on male infertility around the globe. *Reprod Biol Endocrinol.* 2015;13:37-45. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Larsen SH, Wagner G, Heitmann L. Sexual function and obesity. *Int J Obes.* 2007;31(8): 1189-98. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Mukhitdinovna JD. Nay Peritoneal Type of Female Infertility (Mkb-10/N 97.1). *Texas J Med Sci.* 2022;7:77-8. [View at Publisher] [Google Scholar]
- Sepidarkish M, Omani-Samani R, Mansournia MA, Yekaninejad MS, Mardi-Mamaghani A, Vesali S, et al. The casual effect of lifestyle factors on outcomes of assisted reproductive techniques: a protocol study on Iranian infertile couples. *Reprod Health.* 2018;15(1):1-9. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Carson SA, Kallen AN. Diagnosis and management of infertility: a review. *JAMA.* 2021;326(1):65-76. [View at Publisher] [Google Scholar] [DOI] [PMID]
- American Society for Reproductive Medicine. Revised American Society for Reproductive Medicine classification of endometriosis:1996. *Fertility sterility.* 1997;67(5):817-21. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Vinodini N, Chatterjee PK, Amemarsoofi A, Suman V, Pai SR. Evaluation of Liver function with *Moringa oleifera* leaf extract in Cadmium-induced adult Wister albino rats. *Int J plants Animal Environ Sci.* 2014;4(3):103-6. [View at Publisher] [Google Scholar] [DOI]
- Demiray S, Pintado ME, Castro PML. Evaluation of phenolic activities of Turkish medicinal plants; *Tilia argentea*, *crataegi folium* leaves, and *Polygonum bistorta* roots. *World Acad of Sci Eng and Tech.* 2009;3(6):74-9. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Ekor M. The growing use of herbal medicines: issues relating to adverse reactions and challenge in monitoring safety. *Front Pharmacol.* 2014;4:177. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Cajuday LA, Pocsidio GL. Effects of *Moringa oleifera* Lam on the reproduction of mice. *J of Med Plants Res.* 2010;4(12):1115-21. [View at Publisher] [Google Scholar] [DOI]
- Bose CK. Possible role of *Moringa oleifera* Lam. Root in the Epithelia ovarian cancer. *Med Gen Med.* 2007;9(1):26-32. [View at Publisher] [Google Scholar] [PMID]
- Choi JH, Choi KC, Auersperg N, Leung PC. Overexpression of Follicle-Stimulating hormone receptors activates oncogenic pathways in preneoplastic ovarian surface epithelial cells. *J Clin Endocrinol Metab.* 2004;89(11):5508-5516. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Shukla S, Mathur R, Prakash AO. Biochemical and Physiological alterations in female reproductive organs of cyclic rats treated with aqueous extract of *Moringa oleifera* Lam. *Acta Eur Fertil.* 1988;19(4):225-32. [View at Publisher] [Google Scholar] [PMID]
- Stopper H, Schmitt E, Kobras K. Genotoxicity of Phytoestrogens. Mutation Research Foundation and Molecular Mechanism of mutagenesis. 2005;574(1-2):139-55. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Ogunsola OA, Owolabi JO, Fabiyi OS, Nwobi NL, Faluyi B, Akinbola AS. *Moringa* plant parts consumption had effects on reproductive functions in male and female rat models. *J Dent Med Sci.* 2017;16(10):82-6. [View at Publisher] [Google Scholar] [DOI]
- Kohler RE. *Lords of the Fly: Drosophila genetics and the experimental life.* America:University of Chicago Press;1999.P23-9. [View at Publisher] [Google Scholar]
- Mobisson SK, Obembe AO, Ukoh IE, Duru GO. The Role of Hypothalamic Pituitary Gonadal Axis in Aqueous Extract of Cannabis Sativa Induced Male Reproductive Dysfunction of Albino Wistar Rats. *Eur J Pharm and Med Res.* 2018;5(1):71-8. [View at Publisher] [Google Scholar]
- Ehigiator BE, Ozolua RI, Egbogu RO. Evaluation of the sex enhancement and fertility properties of *Waltheria indica* ethanol root extract in *Drosophila melanogaster* (Fruit flies). *Int J of Pharm and Pharm Res.* 2021;3(1):4-8. [View at Publisher] [Google Scholar] [DOI]
- Brooks WH, Daniel KG, Sung SS, Guida WC. Computational validation of the importance of absolute stereochemistry in virtual screening. *J Chem Inf Model.* 2008;48(3):639-45. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Berman HM, Westbrook J, Feng Z, Gilliland G, Bhat TN, Weissig H, Shindyalov IN, et al. The Protein data bank. *Nucleic Acids Res.* 2000;28(1):235-42. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Sastry GM, Adzhigrey M, Day T, Annabhimoju R, Sherman W. Protein and Ligand preparation: Parameters, protocols, and influence on virtual screening enrichments. *J Comput Aided Mol Des.* 2013;27(3):221-34. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Ehigiator BE, Umana IK, Ikem CJ, Iuzegbu C, Ojesebholo RO, Fokunang C. Honey: A remedy for depression. An investigation by experimental validation and molecular docking studies. *Int J Pharm and Pharmaceut Sci.* 2021;3(1):5-15. [View at Publisher] [Google Scholar] [DOI]
- Al-Humesh M, Mahmood AY, Lateef RH. Effects of watery extracts from the root of *Salvadora persia* on the concentration of the sexual hormones and a number of fertility parameters of white mice females. *J Tikrit Uni Agric Sci.* 2012;12(3):62-8. [View at Publisher] [Google Scholar]
- Leone A, Fiorillo G, Criscuolo F, Ravasenghi S, Santogostini L, Fico G, et al. Nutritional characterization of phenolic profiling of *M. oleifera* leaves grown in Chad, Sahrawi refugee camps and Haiti. *Int J Mol Sci.* 2015;16(8):18923-37. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Vongsak B, Sithisam P, Gristanapan W. Simultaneous HPLC Quantitative analysis of active compounds in leaves of *M. oleifera* Lam. *J Chromatogr Sci.* 2014;52(7):641-5. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Atawodi S, Atawodi J, Idakwo G, Pfundstein B, Haubner R, Wurtele G, et al. Evaluation of the polyphenol content and antioxidant properties of methanolic extracts of the leaves, stem, and root barks of *Moringa oleifera* Lam. *J Med Food.* 2010;13(3):710-6. [View at Publisher] [Google Scholar] [DOI] [PMID]
- El-Alfy T, Essat S, Hegazy A, Amer A, Kamel G. Isolation of biologically active constituents of *Moringa oleifera* (Moringaceae) growing in Egypt. *Pharmacogn Mag.* 2011;7(26):109-15. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Muhammad A, Arulselvan P, Cheah P, Abas F, Fakurazi S. Evaluation of the wound healing properties of the tivo aqueous extract of *M. oleifera* Lam on an experimentally induced diabetic animal model. *Drug Des Devel Ther.* 2016;10:1715-30. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Nibret E, Wink M. Trypanocidal and antileukemic effects of the essential oils of *Hagenia abyssinica*, *Leonotis ocyimifolia*, *Moringa stenopetala*, and their main individual constituents. *Phytomedicine.* 2010;17(12):911-20. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Saini R, Sivanesan I, Keum Y. Phytochemicals of *M. oleifera*: a review of their nutritional, therapeutic and industrial significance. *3 Biotech.* 2016;6(2):203. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Verma A, Vijayakumar M, Mathela C, Rao C. In vitro and in vivo antioxidants properties of different fractions of *Moringa oleifera* leaves. *Food Chem Toxicol.* 2009;47(9): 2196-201. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Abd El, Baky H, ElBaroty G. Characterization of Egyptians *M. seed oil* and its bioactivities. *Int J Manag Sci and Bus Res.* 2013;2(7):98-108. [View at Publisher] [Google Scholar]
- Singh BN, Singh BR, Singh RL, Prakash D, Dhakarey R, Upadhyay G. Oxidative DNA damage protective activity, antioxidant and anti quorum sensing potentials of *M. oleifera*. *Food Chem Toxicol.* 2009;47(6):1109-16. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Al-Asmari A, Albalawi S, Athar M, Khan A, Al-Shahrani H, Islam M. *Moringa oleifera* as an anti-cancer agent against breast and colorectal cancer cell lines. *PloS One.* 2015;10(8):e01135814. [View at Publisher] [Google Scholar] [DOI] [PMID]
- Shanker K, Gupta M, Srivastava S, Bawankule D, Pal A, Khanuja S. Determination of active nitrile glycosides in *Moringa oleifera* by reverse phase HPLC. *Food Chem.* 2007;105(1):376-82. [View at Publisher] [Google Scholar] [DOI]
- Sahakitpichan P, Mahidol C, Disadee W, Ruchirawat S, Kanchanapoom T. Unusual glycosides of pyrrole alkaloid and 4'-hydroxyphenylethanamide from leaves of *M. oleifera*. *Phytochemistry.* 2011;72(8):791-5. [View at Publisher] [Google Scholar] [DOI] [PMID]

38. Guevara A, Vargas C, Sakurai H, Fujiwara Y, Hashimoto K, Maoko T, et al. An antitumor promoter from *Moringa oleifera* Lam. *Mutat Res.* 1999;440(2):181-8. [[View at Publisher](#)] [[Google Scholar](#)] [[DOI](#)] [[PMID](#)]
39. Murakami A, Kitazono Y, Jiwajinda S, Koshimizu K, Ohigashi H. Niaziminin, a thiocarbamate from the leaves of *Moringa oleifera*, holds a strict structural requirement for inhibition of tumor-promoter-induced Epstein- Barr Virus activation. *Planta Med.* 1998;64(4):319-23. [[View at Publisher](#)] [[Google Scholar](#)] [[DOI](#)] [[PMID](#)]
40. Kurzer MS, Xu X. Dietary Phytoestrogens. *Annu Rev Nutr.* 1997;17:353-81. [[View at Publisher](#)] [[Google Scholar](#)] [[DOI](#)] [[PMID](#)]
41. Ehigiator BE, Cobhams AE, Adikwu E, Ben IO. Predicting the Possible Mechanism of Aphrodisiac Action of *Pheonix Dactylifera* L. *J Trop Dis.* 2022;10(2):312-20. [[View at Publisher](#)] [[Google Scholar](#)] [[DOI](#)]
42. Herbison AE. Multimodal influence of estrogen upon gonadotropin-releasing hormone neurons. *Endocr Rev.* 1998;19(3):302-30. [[View at Publisher](#)] [[Google Scholar](#)] [[DOI](#)] [[PMID](#)]
43. Konig B, Rauer C, Rosenbaum S, Brandsch C, Eder K, Stangl G. Fasting upregulates PPAR α target genes in the brain and influences pituitary hormone expression in PPAR α dependent manner. *PPAR Res.* 2009;2009:9-12. [[View at Publisher](#)] [[Google Scholar](#)] [[DOI](#)] [[PMID](#)]
44. Abe M, Hojo T, Kozai K, Okuda K. Possible role of insulin-like factor 3 in bovine corpus luteum. *J Vet Med sci.* 2013;75(5):629-32. [[View at Publisher](#)] [[Google Scholar](#)] [[DOI](#)] [[PMID](#)]
45. Kawamura K, Kumagai J, Sudo S, Chun SY, Pisarska M, Morita H, et al. Paracrine regulation of mammalian oocyte maturation and male germ cell survival. *Proc Natl Acad Sci USA.* 2004;101(19):7323-8. [[View at Publisher](#)] [[Google Scholar](#)] [[DOI](#)] [[PMID](#)]
46. Sugimoto Y, Inazumi T, Tsuchiya S. Roles of prostaglandin receptors in female reproduction. *J Biochem.* 2015;157(2):73-80. [[View at Publisher](#)] [[Google Scholar](#)] [[DOI](#)] [[PMID](#)]
47. Grummer RR, Carroll DJ. A review of lipoprotein cholesterol metabolism: Importance to ovarian function. *J Anim Sci.* 1998;66(12):3160-73. [[View at Publisher](#)] [[Google Scholar](#)] [[DOI](#)] [[PMID](#)]

How to Cite:

Ehigiator BE, Iyanyi UL, Mobisson KS, Ukata OI. In vivo and in silico investigation of effects of ethanol extract of *Moringa oleifera* leaves on female fertility, using fruit flies and molecular docking. *JCBR.* 2023;7(2):1-6.



© The author(s)